

Citation: Miller, M. W., and J. R. Fischer. 2016. The First Five (or More) Decades of Chronic Wasting Disease: Lessons for the Five Decades to Come. *Transactions of the North American Wildlife and Natural Resources Conference* 81: in press.

The First Five (or More) Decades of Chronic Wasting Disease: Lessons for the Five Decades to Come

Michael W. Miller

*Colorado Division of Parks and Wildlife
Fort Collins, Colorado*

John R. Fischer

*Southeastern Cooperative Wildlife Disease Study, University of Georgia
Athens, Georgia*

Summary

Chronic wasting disease (CWD), an infectious prion disease of at least five cervid species, has run the gamut from minor scientific curiosity to national crisis since the syndrome's first recognition in the late 1960s. As of April 2016, CWD had been reported in captive and/or free-ranging cervids in the United States (24 states), Canada (three provinces), South Korea, and Norway. With few exceptions (New York and perhaps Minnesota in the US), once in the wild the disease has persisted in reporting jurisdictions in the face of widely varied control attempts. Natural and anthropogenic factors have contributed to the geographic spread and persistence of CWD: Natural factors include prolonged incubation, multiple routes of agent shedding, the agent's environmental persistence, and migratory and dispersal movements of wild cervids; anthropogenic factors include movements of infected live animals (and perhaps infectious tissues and other materials), concentrating susceptible host species, and other artificial wildlife management practices. Many facets of CWD biology and ecology now are well understood, but science informing effective management and control strategies remains comparatively incomplete. Eradicating CWD appears infeasible given its extensive distribution and other epidemiological attributes. Regardless, adaptive approaches for containing foci and reducing infection and transmission rates have shown some promise and deserve further attention. Such pursuits undoubtedly will be more difficult to champion and garner support for in sociopolitical climates ranging from apathetic to combative, particularly when control prescriptions impinge upon or conflict with commercial and sport hunting interests. We believe there are two important motivations for making progress toward sustainable containment and control strategies for CWD in the coming decades: First, data from several sources suggest that heavily-infected cervid populations will not thrive in the long-term. Second, data on CWD prions and experience with other animal prion diseases suggest minimizing human exposure to these agents is prudent.

Chronic Wasting Disease: Lessons Learned from the First Five Decades

Chronic wasting disease (Williams & Young 1980), an infectious prion disease of at least five cervid species (Williams 2005, Norwegian Veterinary Institute 2016), has run the gamut from minor scientific curiosity to national crisis since the syndrome's first recognition in the late 1960s. Moving forward, we believe this wildlife disease merits attention somewhere between

those extremes. Collective experiences and observations made over the last five decades can serve – for better or worse – as a solid foundation for wildlife and animal health professionals to build upon in addressing anticipated challenges posed by CWD in the decades to come. Here we overview what we regard as the key lessons learned over the first five or more decades of North America’s experience with CWD.

Longer Than You Think: Brief History & Known Distribution of Chronic Wasting Disease

That the duration of an outbreak often is underestimated seems perhaps the most important overarching lesson about CWD. Despite its likely occurrence in multiple locations since the 1960s or earlier (Williams & Young 1992, Miller et al. 2000, Wasserberg et al. 2009), many wildlife and animal health professionals, as well as our lay and media publics, perceive CWD as having emerged and spread rapidly only since the early 2000s (e.g., see Saunders et al. 2012). This perception has fostered the broader notion that newly discovered disease foci are truly “new” (i.e., very recent) occurrences. To the contrary, given imperfect surveillance approaches, incomplete or inaccurate knowledge about local exposure risks, and the insidious progression of an outbreak in its early stages, the first case *detected* in a locale is rarely *THE* first case that has occurred. Consequently, on further investigation “new” foci tend to have larger spatial dimensions and higher prevalence than expected, thereby perpetuating misconceptions about the speed of “spread.” This lesson has been illustrated by experiences in Colorado and Wyoming (Miller et al. 2000), in Saskatchewan (Bollinger et al. 2004, Argue et al. 2007), in Wisconsin (Wasserberg et al. 2009, Holsman et al. 2010), and most recently in Arkansas where expanded surveillance disclosed 79 additional cases within two month after their “first” case was diagnosed in February 2016 (Arkansas Game & Fish Commission 2016).

Chronic wasting disease history (Table 1) remains incompletely documented. The “chronic wasting” syndrome first was recognized in captive mule deer (*Odocoileus hemionus*) deer held for research in Colorado in the 1960s (Williams & Young 1980), but unrecognized cases could have occurred in Colorado or elsewhere before that time (Williams & Miller 2003). Clinical cases also were recognized in captive mule deer in the Denver and Toronto zoos in the 1970s, and in captive Rocky Mountain elk (*Cervus elaphus nelsoni*) in research and zoological collections in Colorado and Wyoming (Williams & Young 1992, Dubé et al. 2006). Undocumented involvement of other private collections or menageries during the 1960s and 1970s seems likely. Within little more than the first two decades after its characterization as a transmissible spongiform encephalopathy (Williams & Young 1980), CWD cases were diagnosed in wild mule deer, white-tailed deer (*Odocoileus virginianus*) and elk in northeastern Colorado and southeastern Wyoming (1980s–1990s), in commercial captive elk facilities in Saskatchewan (1996) and in South Dakota (1997), in commercial captive white-tailed deer facilities in several jurisdiction (2001–2002), and eventually in moose (*Alces alces*; Williams & Young 1992, Spraker et al. 1997, Miller et al. 2000, Williams et al. 2002, Williams & Miller 2003, Williams 2005). Cases from what have become recognized as large foci in Saskatchewan-Alberta and Wisconsin-Illinois also were first detected in the early 2000s (Williams et al. 2002). As of April 2016, cases of CWD had been reported in captive and/or free-ranging cervids in 24 US states (75 captive herds in 16 states and free-ranging cervids in 22 states), three Canadian provinces (including Ontario’s Toronto Zoo in the 1970s), and South Korea (Fig. 1; Williams & Young 1992, Williams et al. 2002, U.S. Geological Survey 2016; R. Pritchard pers. comm.). In addition, at the time of this writing a single case in a free-ranging reindeer (*Rangifer tarandus*) in

Norway in March 2016 had just been reported (Norwegian Veterinary Institute 2016). Based on experience to date, the true geographic distribution of CWD likely remains underestimated.

Two Good Stories: The Drivers of Chronic Wasting Disease “Spread”

A second overarching lesson – a corollary to the first – is that new CWD foci often can be explained by two or more equally plausible (and equally undeniable) “origin stories.” Distorted temporal perceptions on the likely timing of introduction underlie the plurality of origin stories, as do sociopolitical motivations to deflect or lay blame elsewhere when “new” cases arise. But perhaps most pervasive is the lack of complete information on contributory events, particularly for outbreaks involving free-ranging cervids. Although the lack of a singular explanation can be dissatisfying, failing to consider plausible alternative timelines and exposure sources may be more problematic when disease prevention and control efforts are misinformed or misled. For example, the widely held belief that all CWD occurrences can be traced back to a single Colorado research facility has precluded wildlife and animal health professionals from considering that some outbreaks may be arising from unrecognized exposure events that occur repeatedly over time (e.g., Williams & Miller 2003, Greenlee et al. 2015). The recent Norwegian reindeer case may stimulate broader thinking.

In fact, both natural and anthropogenic factors have contributed to the geographic spread and persistence of CWD over the last five decades. Regardless of the ultimate origin, much of the geographic “spread” of CWD appears attributable to natural movements in some jurisdictions: Wyoming, for example, has only one private game farm and consequently commercial enterprise is unlikely to have driven the widespread distribution there. Alternatively, the role of commercial elk operations in CWD outbreaks in Saskatchewan and South Korea was well-documented (Williams et al. 2002, Kim et al. 2005, Argue et al. 2007), with inadvertent spill-over also giving rise to a large free-ranging focus spanning the Saskatchewan-Alberta shared border (Bollinger et al. 2004). In Colorado, a combination of natural and anthropogenic factors likely contributed in different measures to separate outbreaks along the Front Range and on the Western Slope.

Natural factors contributing to persistence and geographic spread include prolonged incubation, multiple routes of agent shedding, the agent’s environmental persistence, and movements of free-ranging cervids. Infected cervids likely shed prions for most of the disease course (Tamgüney et al. 2009, Henderson et al. 2015), thus affording ample opportunities for transmission within and among social groups. Migration movements also have potential for contributing to longer-distance jumps in distribution. Because infectivity can be harbored in some environments for an extended time, transmission occurs on overlapping ranges even in the absence of direct interactions between infected and uninfected animals. Indirect transmission also increases the likelihood of interspecies transmission.

The primary anthropogenic factor identified in the dissemination of CWD is human-facilitated movement of live animals (Williams & Miller 2003), and to date this is the only confirmed contributing activity linked to CWD’s spread between distant locations. These animal movements typically are fostered by other highly artificial wildlife management activities, such as captive wildlife propagation and high-fenced shooting enclosures (Fischer and Davidson, 2005). Although spared from implication thus far, translocating free-ranging cervids from an infected source also would present a similar risk for spreading CWD. Local wildlife may be exposed to CWD if infected captive animals escape, or if there is ingress/egress of free-ranging cervids with exposure to infected captive animals or to contaminated environments. Fence-line

contact offers another opportunity for direct transmission. (We note that these transmission opportunities are a two-way street, i.e., CWD can move in either direction between captive and wild cervids.) Other possible modes for the anthropogenic spread of CWD include transport of infected carcasses, products manufactured or contaminated with prion-laden deer or elk urine, saliva, or feces, and movement of hay or grain crops contaminated with the CWD agent. None of these has been documented in the field, although proof of concept has been demonstrated experimentally.

In addition, other anthropogenic factors can substantially increase the likelihood of establishing, maintaining, and disseminating CWD and other diseases in free-ranging wildlife. In particular, artificial management activities, such as wildlife baiting and feeding or other practices that congregate normally dispersed wild animals, enhance pathogen transmission opportunities (Fischer and Davidson, 2005).

Things We Now Know: Chronic Wasting Disease Biology & Ecology

Many facets of CWD biology and ecology that were mysteries even into the early 2000s (e.g., Williams et al. 2002) now are well understood. For example, notable advances have been made in diagnostics and in our understanding of transmission routes and host factors modulating disease progression that have application in CWD detection and control. These and other advances have been reviewed thoroughly elsewhere (e.g., Williams 2005, Sigurdson 2008, Smith et al. 2011, Saunders et al. 2012, Haley & Hoover 2015). We offer here only a brief synthesis of findings most relevant to detection and control, referring interested readers to the aforementioned reviews and numerous original papers referenced therein for greater details on and sources of specific points highlighted in our synopsis:

Chronic wasting disease appears to be caused by one or more strains of infectious prions. Although the ultimate historical origin never will be known with certainty, we regard exposure of native cervids to the sheep scrapie agent at one or more times and locations as a parsimonious explanation. Regardless of their origin(s), sustained outbreaks now occur as large and small foci and in captive wildlife facilities (Fig. 1). Natural cases of CWD have occurred in five host species: mule deer, white-tailed deer, Rocky Mountain elk, moose, and reindeer/caribou. No immunity, recovery, or absolute resistance to infection has been documented in any of the susceptible species. However, natural variation in the host gene encoding for cellular prion protein does modulate disease progression, thereby extending survival times and perhaps lowering infection probabilities for “relatively resistant” genotypes. The disease course typically is measured in years. Clinical signs – altered behavior initially, with body condition declining much later – become progressively apparent relatively late in the disease course. Infection can be detected in carcasses, as well as in live animals, and diagnostic tests become increasingly reliable in individual animals as the disease progresses. Chronic wasting disease is infectious. Infected individuals shed prions from several routes during most of the disease course, exposing others either directly or through contamination of shared resources or environments. Shed prions can persist for years in the environment, and their binding to soil elements (e.g., clay) enhances persistence and infectivity. The uncoupling of transmission from the immediate presence of infected animals greatly complicates CWD control.

Looking Hard–Hardly Looking: Detecting Chronic Wasting Disease

A third key lesson relates to the difficulty in detecting CWD foci in captive and wild settings despite the considerable effort expended. Most North American jurisdictions have, at

least for a time since the early 2000s, engaged in extensive if not intensive surveillance to seek out such foci. Although all of these efforts were well-intentioned, many were too flawed or too short-lived to provide reliable information on disease absence. We briefly review common shortcomings of CWD surveillance as widely practiced to provide a basis for improving the efficiency and effectiveness of future efforts.

Preferred approaches for seeking out new foci (termed “surveillance” here) differ from approaches for following epidemic trends over time (“monitoring;” concepts reviewed in greater detail by Samuel et al. 2003). We recommend that CWD surveillance be an ongoing activity in jurisdictions or areas where foci have not been detected previously; monitoring may be a more episodic undertaking (e.g., at multi-year intervals) where support resources are limited because infection rates tend to change slowly compared to more conventional infectious diseases. Regardless of the purpose, CWD surveillance and monitoring should be undertaken at biologically relevant spatial scales and inferences drawn only in the appropriate spatial context in view of the highly patchy distribution of CWD in wild cervids. In our experience, statements exhorting that examination of a few hundred (or even a few thousand) harvested animals has proven a jurisdiction’s freedom from CWD rarely are supported by the data in hand.

For surveillance in free-ranging settings, targeting sample sources known to have a relatively high probability of infection in endemic areas (e.g., clinical “suspects,” vehicle- or predator-killed adult animals) can be a more cost-effective approach (Miller et al. 2000, Samuel et al. 2003, Walsh & Miller 2010). The effectiveness of so-called “targeted surveillance” assumes relatively even sampling effort over the geographic area of inference. However, this approach does have limitations: For example, clinical disease may not be observed in remote areas, vehicle-kills do not occur in roadless areas, and predator kills may be consumed before sampling can occur. In addition to clinical targeting, spatial targeting via risk-based assessments also may enhance the effectiveness and efficiency of CWD surveillance (Bollinger et al. 2004, Rees et al. 2012, Norbert et al. 2016).

For monitoring, random sampling (e.g., from harvested animals) provides relatively unbiased estimates of infection rates (Samuel et al. 2003). Comparisons over time or between locations should be based on a common denominator (e.g., harvested males aged 2 years or older) to assure that reliable inferences are drawn. Where available, data from lethal and nonlethal sampling can be combined for analysis provided sources are equivalent (e.g., Geremia et al. 2015). Because foci emerge and grow so slowly, infection rates may be remarkably high on first detection when jurisdictions rely on random sampling for surveillance. Moreover, CWD tends to be unevenly distributed in the wild. The notion that a survey sample of 300 assures 95% probability of detecting at least one case where prevalence $\geq 1\%$ assumes infection is evenly distributed at that rate throughout the entire target population (Samuel et al. 2003). However, CWD distribution typically is uneven within an affected population, and the target population itself often is distributed unevenly.

Toward a Sustained & Sustainable Effort to Control Chronic Wasting Disease

The final overarching lessons learned over the past five decades relate to how wildlife and animal health professionals should (and probably should not) approach the control of CWD. In contrast to advances in our understanding of CWD biology and ecology, the available science informing effective management and control strategies remains relatively incomplete. However, recent insights and modest strides seem to offer a path forward. It follows that adaptive

approaches for containing CWD foci and reducing infection and transmission rates deserve further attention.

Eradicating CWD from North America appears infeasible given its extensive distribution and other epidemiological attributes. With few exceptions –the small foci in New York (New York Department of Environmental Conservation 2015) and perhaps Minnesota (Minnesota Department of Natural Resources 2014) – CWD in free-ranging cervids has persisted in reporting jurisdictions in the face of widely varied control attempts. Faced with the dim prospects for eradication on scales large or small, some affected jurisdictions now seem to have abandoned any further consideration of disease management and some have effectively dismantled surveillance and monitoring as well. In light of myriad wildlife conservation needs and ever-dwindling resources we appreciate the allure but believe this to be myopic. Instead, we strongly encourage affected jurisdictions to redouble efforts to collectively foster and develop sustained and sustainable approaches for CWD surveillance, monitoring, and control.

In contrast to the apparent success in eliminating New York's small free-ranging focus, well-publicized early attempts to control CWD in Colorado and Wisconsin yielded little evidence of progress and thus gave initial appearances of failure (e.g., Conner et al. 2007, Holsman et al. 2010). In recent years, however, evidence from several jurisdictions' control attempts applied across different geographic scales suggest that combinations of intensive deer removal focused around case clusters and more sustained suppression of the affected herd or population may offer some measure of effective disease suppression. A sustained culling program underway since 2003 has stabilized prevalence in northern Illinois white-tailed deer as compared to increasing trends in southern Wisconsin where disease control was suspended in 2007 (Manjerovic et al. 2014). Similar divergence in prevalence among white-tailed and mule deer harvested in Alberta and Saskatchewan (Norbert et al. 2016) may reflect the relative effectiveness of disease suppression approaches in Alberta (Pybus 2012), but also could be an artifact of more recent epizootic emergence in Alberta. In northcentral Colorado, a combination of focal culling (Conner et al. 2007) and broader, harvest-mediated population reduction (~25%; Colorado Parks & Wildlife, unpubl. data) in the early 2000s appears likely to have contributed to reduced prevalence (Geremia et al. 2015), whereas estimated prevalence in other Colorado mule deer herds has increased since 2002 (Colorado Parks & Wildlife, unpubl. data).

One of the most common flaws in CWD control efforts to date has been initial underestimation of the affected area (often based on inadequate surveillance and erroneous assumptions about how long disease has been present). The outcome then gave the appearance that the control attempt had "failed" when in fact the approach was biologically sound but the application was either too small (spatially) or too short-lived. It follows that acquiring reliable distribution and prevalence data early in the planning and execution may improve the apparent efficacy of future CWD control efforts. To this end, we encourage jurisdictions to consider and set realistic disease control objectives and to use adaptive management approaches that incorporate existing and prospective field data to refine disease control objectives.

In addition to adopting and adaptively assessing approaches for stabilizing or suppressing CWD epizootics, we encourage jurisdictions to consider how recent trends in cervid management may be contributing to disease emergence. Modeling suggests harvest-based control of CWD may be most effective when focused on male deer (Jennelle et al. 2014, Potapov et al. 2016), perhaps because infection rates among adult male deer tend to be higher than among adult females (Miller et al. 2000, Grear et al 2006, Rees et al. 2012). Conversely, then, harvest strategies intended to increase male:female ratios or adult male age structure could inadvertently

facilitate CWD persistence. This may explain why in Colorado, for example, the dramatic increases in prevalence observed since 2002 in several affected mule deer herds coincide with statewide changes in harvest strategies that have increased male:female ratios over the same period (Bergman et al. 2011). Given the potential for unintended consequences, we encourage broader critical assessment of how this and other harvest strategies (e.g., season timing, baiting, “quality deer management”) may be affecting CWD dynamics.

Such pursuits undoubtedly will be more difficult to champion and garner support for in sociopolitical climates ranging from apathetic to combative, particularly when control prescriptions impinge upon or conflict with commercial and sport hunting interests. The human dimensions of managing wildlife diseases in general and CWD in particular present a substantial challenge for those determining the management objectives and actions. For example, surveys of hunters and landowners in Wisconsin identified factors that contributed to hunter opposition to the state’s CWD management plan including: opposition to deer population goals (initially zero), conflicts with traditions and consumption norms, uncertainty about the likelihood of success, questions about agency credibility, and no sense of urgency (Holsman et al. 2010).

We believe there are two important motivations for responsible wildlife managers to make progress toward sustainable containment and control strategies for CWD in the coming decades: First, data from several sources suggest that heavily-infected cervid populations will not thrive in the long-term (Miller et al. 2000, 2008, Almberg et al. 2011, Edmunds 2013, Monello et al. 2014, Williams et al. 2014, DeVivo 2015). Second, data on CWD prions and experience with other animal prion diseases suggest minimizing human exposure to these agents would be prudent (Raymond et al. 2000, Belay et al. 2004, Saunders et al. 2012, Cassard et al. 2014).

References

- Almberg, E. S., P. C. Cross, C. J. Johnson, D. M. Heisey, and B. J. Richards. 2011. Modeling Routes of Chronic Wasting Disease Transmission: Environmental Prion Persistence Promotes Deer Population Decline and Extinction. *PLoS ONE* 6:e19896. doi:10.1371/journal.pone.0019896
- Argue, C. K., C. Ribble, V. W. Lees, J. McLane, and A. Balachandran. 2007. Epidemiology of an Outbreak of Chronic Wasting Disease on Elk Farms in Saskatchewan. *Canadian Veterinary Journal* 48:1241–1248.
- Arkansas Game and Fish Commission. 2016. CWD Jasper Public Meeting Summary. Thursday, April 7 2016. Web 14 April 2016. <http://www.agfc.com/hunting/Documents/CWD-JasperMeetingSummary160407.pdf>
- Belay, E. D., R. A. Maddox, E. S. Williams, M. W. Miller, P. Gambetti, and L. B. Schonberger. 2004. Chronic Wasting Disease and Potential Transmission to Humans. *Emerging Infectious Diseases* 10(6): 977–984.
- Bergman, E. J., B. E. Watkins, C. J. Bishop, P. M. Lukacs, and M. Lloyd. 2011. Biological and Socio-Economic Effects of Statewide Limitation of Deer Licenses in Colorado. *The Journal of Wildlife Management* 75(6):1443–1452; 2011; DOI: 10.1002/jwmg.168
- Bollinger, T., P. Caley, E. Merrill, F. Messier, M.W. Miller, M.D. Samuel, and E. Vanopdenbosch. 2004. Chronic Wasting Disease in Canadian Wildlife: An Expert Opinion on the Epidemiology and Risks to Wild Deer. Prepared by Expert Scientific Panel on Chronic Wasting Disease. July 2004. Canadian Cooperative Wildlife Health Centre, Western College of Veterinary Medicine, Saskatoon, Sask. Available from http://www.cwhc-rcsf.ca/docs/technical_reports/CWD_Expert_Report_Final_20040804.pdf [Accessed 14 April 2016]
- Cassard, H., J.-M. Torres, C. Lacroux, J.-Y. Douet, S. L. Benestad, F. Lantier, S. Lugin, I. Lantier, P. Costes, N. Aron, F. Reine, L. Herzog, J.-C. Espinosa, V. Beringue, and O. Andréoletti. 2014. Evidence for Zoonotic Potential of Ovine Scrapie Prions. *Nature Communications* 5:5821. doi: 10.1038/ncomms5821.
- Conner, M. M., M. W. Miller, M. R. Ebinger, and K. P. Burnham. 2007. A Meta-BACI Approach for Evaluating Management Intervention on Chronic Wasting Disease in Mule Deer. *Ecological Applications* 17 (1): 140–153.
- DeVivo, M. T. 2015. Chronic Wasting Disease Ecology and Epidemiology of Mule Deer in Wyoming. Ph.D. Dissertation. University of Wyoming. 154 pp.

- Dubé, C., K. G. Mehren, I. K. Barker, B. L. Peart, and A. Balachandran. 2006. Retrospective Investigation of Chronic Wasting Disease of Cervids at the Toronto Zoo, 1973–2003. *Canadian Veterinary Journal* 47:1185–1193.
- Edmunds, D. R. 2013. Chronic Wasting Disease Ecology and Epidemiology of White-tailed Deer in Wyoming. Ph.D. Dissertation. University of Wyoming. 200 pp.
- Fischer, J. R., and W. R. Davidson. 2005. Reducing Risk Factors for Disease Problems Involving Wildlife. *Transactions of the North American Wildlife and Natural Resources Conference* 70:289-309.
- Geremia, C., M. W. Miller, J. A. Hoeting, M. F. Antolin, and N. T. Hobbs. 2015. Bayesian Modeling of Prion Disease Dynamics in Mule Deer Using Population Monitoring and Capture-Recapture Data. *PLoS ONE* 10(10): e0140687. doi:10.1371/journal.pone.0140687
- Greear, D. A., M. D. Samuel, J. A. Langenberg, and D. Keane. 2006. Demographic Patterns and Harvest Vulnerability of Chronic Wasting Disease Infected White-tailed Deer in Wisconsin. *Journal of Wildlife Management* 70: 546–553. doi:10.2193/0022-541X(2006)70[546:DPAHVO]2.0.CO;2
- Greenlee, J., S. J. Moore, J. Smith, M. H. W. Greenlee, and R. Kunkle. 2015. Scrapie Transmits to White-Tailed Deer by the Oral Route and has a Molecular Profile Similar to Chronic Wasting Disease and Distinct from the Scrapie Inoculum. *PRION* 9: S62–S62 (Supplement 1, Meeting Abstract P.97, published 24 April 2015).
- Haley, N. J., and E. A. Hoover. 2015. Chronic Wasting Disease of Cervids: Current Knowledge and Future Perspectives. *Annual Review of Animal Biosciences* 3:305–325. doi: 10.1146/annurev-animal-022114-111001. Epub 2014 Oct 2.
- Henderson, D. M., N. D. Denkers, C. E. Hoover, N. Garbino, C. K. Mathiason, and E. A. Hoover. 2015. Longitudinal Detection of Prion Shedding in Saliva and Urine by Chronic Wasting Disease-Infected Deer by Real-Time Quaking-Induced Conversion. *Journal of Virology* 89:9338 –9347. doi:10.1128/JVI.01118-15.
- Holsman, R. H., J. Petchenik, and E. E. Cooney. 2010. CWD After “The Fire:” Six Reasons Why Hunters Resisted Wisconsin’s Eradication Effort. *Human Dimensions of Wildlife* 15(3):180–193.
- Jennelle, C. S., V. Henaux, G. Wasserberg, B. Thiagarajan, R. E. Rolley, and M. D. Samuel. 2014. Transmission of Chronic Wasting Disease in Wisconsin White-Tailed Deer: Implications for Disease Spread and Management. *PLoS ONE* 9(3): e91043. doi:10.1371/journal.pone.0091043
- Kim, T.-Y., H.-J. Shon, Y.-S. Joo, U.-K. Mun, K.-S. Kang, and Y.-S. Lee. 2005. Additional Cases of Chronic Wasting Disease in Imported Deer in Korea. *Journal of Veterinary Medical Science* 67(8):753–759.
- Manjerovic, M. B., M. L. Green, N. Mateus-Pinilla, and J. Novakofski. 2014. The Importance of Localized Culling in Stabilizing Chronic Wasting Disease Prevalence in White-tailed Deer Populations. *Preventive Veterinary Medicine* 113:139–145. doi: 10.1016/j.prevetmed.2013.09.011
- Miller, M. W., E. S. Williams, C. W. McCarty, T. R. Spraker, T. J. Kreeger, C. T. Larsen, and E. T. Thorne. 2000. Epizootiology of Chronic Wasting Disease in Free-ranging Cervids in Colorado and Wyoming. *Journal of Wildlife Diseases* 38: 676–690.
- Miller, M. W., H. M. Swanson, L. L. Wolfe, F. G. Quartarone, S. L. Huwer, C. H. Southwick, and P. M. Lukacs. 2008. Lions and Prions and Deer Demise. *PLoS ONE* 3(12): e4019. doi:10.1371/journal.pone.0004019.
- Minnesota Department of Natural Resources. 2014. Managing Chronic Wasting Disease. Web 13 April 2016. <http://www.dnr.state.mn.us/hunting/deer/cwd/index.html>
- Monello, R. J., J. G. Powers, N. T. Hobbs, T. R. Spraker, M. K. Watry, and M. A. Wild. 2014. Survival and Population Growth of a Free-ranging Elk Population with a Long History of Exposure to Chronic Wasting Disease. *The Journal of Wildlife Management* 78(2): 214–223.
- New York Department of Environmental Conservation. 2015. Status of CWD. Web 7 April 2016. <http://www.dec.ny.gov/animals/33220.html>
- Norbert, B. R., E. H. Merrill, M. J. Pybus, T. K. Bollinger, and Y. T. Hwang. 2016. Landscape Connectivity Predicts Chronic Wasting Disease Risk in Canada. *Journal of Applied Ecology* (in press).
- Norwegian Veterinary Institute. 2016. The First Detection of Chronic Wasting Disease (CWD) in Europe. Web. 4 April 2016. (<http://www.vetinst.no/eng/Highlights/The-first-detection-of-Chronic-Wasting-Disease-CWD-in-Europe>)
- Potapov, A., E. Merrill, M. Pybus, and M. A. Lewis. 2016. Chronic Wasting Disease: Transmission Mechanisms and the Possibility of Harvest Management. *PLoS ONE* 11(3): e0151039. doi:10.1371/journal.pone.0151039

- Pybus, M. J. 2012. CWD Program Review 2012. Alberta Sustainable Resource Development, Fish and Wildlife Division. Web 17 March 2016. <http://aep.alberta.ca/fish-wildlife/wildlife-diseases/chronic-wasting-disease/documents/CWD-ProgramReview-May-2012.pdf>
- Raymond, G. J., A. Bossers, L. D. Raymond, K. I. O'Rourke, L. E. McHolland, P. K. Bryant III, M. W. Miller, E. S. Williams, M. Smits, and B. Caughey. 2000. Evidence of a Molecular Barrier Limiting Susceptibility of Humans, Cattle, and Sheep to Chronic Wasting Disease. *The EMBO Journal* 19(17): 4425–4430.
- Rees, E. E., E. H. Merrill, T. K. Bollinger, Y. T. Hwang, M. J. Pybus, and D. W. Coltman. 2012. Targeting the Detection of Chronic Wasting Disease Using the Hunter Harvest during Early Phases of an Outbreak in Saskatchewan, Canada. *Preventive Veterinary Medicine* 104:149–159. doi: 10.1016/j.prevetmed.2011.10.016
- Samuel, M. D., D. O. Joly, M. A. Wild, S. D. Wright, D. L. Otis, R. W. Werge, and M. W. Miller. 2003. Surveillance Strategies for Detecting Chronic Wasting Disease in Free-ranging Deer and Elk. Results of a CWD Surveillance Workshop. USGS, BRD, National Wildlife Health Center, Madison, Wisconsin. 41 pp. http://www.nwhc.usgs.gov/research/chronic_wasting/CWD_Surveillance_Strategies.pdf
- Saunders, S. E., S. L. Bartelt-Hunt, and J. C. Bartz. 2012. Occurrence, Transmission, and Zoonotic Potential of Chronic Wasting Disease. *Emerging Infectious Diseases* 18:369–376. DOI: <http://dx.doi.org/10.3201/eid1803.110685>
- Sigurdson, C.J. 2008. A Prion Disease of Cervids: Chronic Wasting Disease. *Veterinary Research* 39:41. <http://dx.doi.org/10.1051/vetres:2008018>
- Smith, C. B., C. J. Booth, and J. A. Pedersen. 2011. Fate of Prions in Soil: A Review. *Journal of Environmental Quality* 40:449–61. <http://dx.doi.org/10.2134/jeq2010.0412>
- Spraker, T. R., M. W. Miller, E. S. Williams, D. M. Getzy, W. J. Adrian, G. G. Schoonveld, R. A. Spowart, K. I. O'Rourke, J. M. Miller, and P. A. Merz. 1997. Spongiform Encephalopathy in Free-Ranging Mule Deer (*Odocoileus Hemionus*), White-Tailed Deer (*Odocoileus Virginianus*), and Rocky Mountain Elk (*Cervus Elaphus Nelsoni*) in Northcentral Colorado. *Journal of Wildlife Diseases* 33: 1–6.
- Tamgüney, G., M. W. Miller, L. L. Wolfe, T. M. Sirochman, D. V. Glidden, C. Palmer, A. Lemus, S. J. DeArmond, and S. B. Prusiner. 2009. Asymptomatic Deer Excrete Infectious Prions in Faeces. *Nature* 461 (7263): 529–U90. DOI: 10.1038/nature08289
- United States Geological Survey. 2016. Chronic Wasting Disease (CWD). National Wildlife Health Center. Web. 1 April 2016. (http://www.nwhc.usgs.gov/disease_information/chronic_wasting_disease/)
- Walsh, D. P., and M. W. Miller. 2010. A Weighted Surveillance Approach for Detecting Chronic Wasting Disease Foci. *Journal of Wildlife Diseases* 46(1): 118–135.
- Wasserberg, G., E. E. Osnas, R. E. Rolley, and M. D. Samuel. 2009. Host Culling As an Adaptive Management Tool for Chronic Wasting Disease in White-Tailed Deer: A Modeling Study. *Journal of Applied Ecology* 46:457–466. doi: 10.1111/j.1365-2664.2008.01576.x
- Williams, A. L., T. J. Kreeger, and B. A. Schumaker. 2014. Chronic Wasting Disease Model of Genetic Selection Favoring Prolonged Survival in Rocky Mountain Elk (*Cervus elaphus*). *Ecosphere* 5(5):60. <http://dx.doi.org/10.1890/ES14-00013.1>
- Williams, E. S. 2005. Chronic Wasting Disease. *Veterinary Pathology* 42: 530–549.
- Williams, E. S., and M. W. Miller. 2003. Transmissible Spongiform Encephalopathies in Non-Domestic Animals: Origin, Transmission, and Risk Factors. In *Risk Analysis of Prion Diseases in Animals*. C. I. Lasmézas and D. B. Adams, (eds.). *Revue scientifique et technique Office international des Epizooties* 22: 145–156.
- Williams, E. S., and S. Young. 1980. Chronic Wasting Disease of Captive Mule Deer: A Spongiform Encephalopathy. *Journal of Wildlife Diseases* 16: 89–98.
- Williams, E. S., and S. Young. 1992. Spongiform Encephalopathies of Cervidae. In *Transmissible Spongiform Encephalopathies of Animals*, R. Bradley and D. Mathews (eds.). *Revue scientifique et technique Office international des Epizooties* 11: 551–567.
- Williams, E. S., M. W. Miller, and E. T. Thorne. 2002. Chronic Wasting Disease: Implications and Challenges for Wildlife Managers. *Transactions of the North American Wildlife and Natural Resources Conference* 67: 87–103.

Table 1. An abbreviated timeline of select chronic wasting disease events. Adopted and from Williams et al. (2002); updated and compiled by K. Niedringhaus, Southeastern Cooperative Wildlife Disease Study. See text for additional details and references.

Year	Events
1967	Wasting syndrome observed in captive mule deer at a Colorado wildlife research facility
1975–81	Wasting syndrome observed in Toronto Zoo mule deer that came from the Denver Zoo
1978	“Chronic wasting disease” (CWD) diagnosed as transmissible spongiform encephalopathy (TSE)
1979	Recognized in captive mule deer at Wyoming wildlife research facility
1981	Detected in wild elk in Colorado
1985	Detected in wild mule deer in Colorado & Wyoming
1996	Detected in a captive elk farm in Saskatchewan; 38 other linked farms eventually found positive
1997	Detected in captive elk facilities in South Dakota
1998	- Detected in captive elk facilities in Montana & Oklahoma <i>Model Program for Surveillance, Control, & Eradication of CWD in Domestic Elk</i> presented at US Animal Health Association to establish monitoring & control standards
1999	World Health Organization indicates no evidence CWD is transmissible to humans, but advises that exposure should be avoided nonetheless
2000	- Detected in wild mule deer in Nebraska & Saskatchewan - Research: molecular studies compare host ranges for CWD, scrapie, & bovine spongiform encephalopathy prions; environmental contamination & subclinical infection contribute to transmission; prevalence estimates in wild populations in Colorado & Wyoming
2001	- Detected in captive elk in Kansas - Detected in captive elk in South Korea imported from Saskatchewan - Detected in wild white-tailed deer in South Dakota - USDA declares CWD emergency in captive elk; funds available for disease control
2002	- Detected in captive elk in Minnesota & captive white-tailed deer in Alberta - Detected in wild & captive white-tailed deer in Wisconsin - Detected in wild white-tailed deer in Illinois, mule deer in New Mexico, elk in South Dakota - Joint CWD Task Force of USDA/DOI/States/Universities develops <i>Plan for Assisting States, Federal Agencies & Tribes in Managing CWD in Wild & Captive Cervids</i> (National CWD Plan) - Colorado establishes guidelines to minimize transport of high risk carcass materials 1st International CWD Symposium (Denver, Colorado) - Research: tonsil biopsy as a live animal test; improved high-throughput diagnostics
2003	- Detected in wild mule deer in Utah - APHIS funds available for CWD work in captive & wild cervids (through 2011) - USDA publishes Proposed Rule for CWD herd certification & interstate shipping program (HCP) to eradicate CWD from captive deer & elk - Research: horizontal transmission of CWD likely important in CWD epidemiology
2004	- Detected in wild elk in New Mexico - National CWD Plan progress report published & new priorities discussed - Research: environmental sources, decomposed carcasses can contribute to transmission
2005	- Detected in captive & wild white-tailed deer in New York - Detected in wild mule deer in Alberta, moose in Colorado, white-tailed deer in West Virginia
2006	- Detected in captive white-tailed deer in Minnesota - Detected in wild white-tailed deer in Kansas - USDA publishes CWD HCP Final Rule – never implemented

	• Research: prions in muscles of infected deer; transmitted in saliva & blood
2007	• Research: prions in environment more infective in particular (clay) soil types
2008	• Detected in captive deer in Michigan • Detected in wild elk in Saskatchewan, moose in Wyoming • Research: CWD may be a plausible explanation for local deer population declines in Colorado;
2009	• APHIS plans to withdraw 2006 CWD Final Rule, issue a new rule based on 2006 rule & 2009 proposed rule • Research: prions shed in feces from deer in early stages of CWD; prions in urine & saliva
2010	• Detected in captive white-tailed deer in Missouri • Detected in wild white-tailed deer in North Dakota & Virginia
2011	• Detected in wild white-tailed deer in Maryland • Detected in wild white-tailed deer in Minnesota • Severe reduction of USDA funds for CWD work
2012	• Detected in captive white-tailed deer in Iowa & Pennsylvania • Detected in wild white-tailed deer in Missouri • Detected in wild mule deer in west Texas • APHIS Interim Final Rule for CWD Herd Certification & Interstate Movement & CWD Program Standards published • Research: possible link between scrapie & CWD
2013	• Detected in wild white-tailed deer in Pennsylvania
2014	• Detected in captive deer in Ohio • CWD Program Standards revised • APHIS CWD Final Rule implemented • Research: plants may play role in CWD transmission & environmental maintenance; experimental aerosol transmission in white-tailed deer
2015	• Detected in wild white-tailed deer in Michigan • Detected in captive white-tailed deer in Texas • Research: plants can bind prions superficially & uptake prions from contaminated soil
2016	• Detected in wild elk & white-tailed deer in Arkansas • Detected in a wild reindeer in Norway




Figure 1. Current known distribution of chronic wasting disease (CWD). In addition to North America, cases have been reported in South Korea (captive only) and Norway (free-ranging only). North America map from U.S. Geological Survey (2016); global maps from Wikipedia.